



A Study On Psycho – Social Impact Of Perimenopause And Menopause Among The Working Women Of Coimbatore District – A Social Work Perspective

***Rasha Ashraf**

**Assistant Professor, Department of Social Work, Nehru Arts and Science College, Coimbatore, Tamil Nadu, India*

Received: 15/05/2026

Revised: 12/08/2026

Accepted: 28/08/2026

Published: 22/09/2026

ABSTRACT

The present study delves into the psychosocial impact of Perimenopause and Menopause on the mental health of working women in Coimbatore District, Tamil Nadu. It explores the effects of hormonal fluctuations, psychosocial factors, coping strategies, and support systems on mental health during this stage. Employing an explanatory research design, it aims to understand causal relationships and complex interactions, elucidating reasons behind mental health changes. The qualitative method of case study was used for data collection, involving a sample of 10 working women under 50 experiencing menopause or its transitional phase. Probability sampling was employed to ensure generalizability. The Findings reveal a correlation between hormonal changes (e.g., decreased Estrogen levels) and symptoms like Depression, Anxiety, mood swings, and cognitive changes. Psychosocial stressors from work, family, and societal attitudes towards aging significantly impact mental health during this transition. The study underscores the need for tailored interventions and support programs to address women's specific needs during perimenopause and menopause, enhancing mental health outcomes and overall well-being. The study concludes with the importance of understanding the multifaceted nature of menopausal experiences, considering both biological and psychosocial factors. By recognizing the challenges faced by working women during this transition, appropriate interventions can be developed to support their mental health effectively as well as it contributes valuable insights to the development of holistic approaches aimed at promoting well-being and resilience among women navigating perimenopause and menopause in Coimbatore District.

Key Words –*Perimenopause, Menopause, Transitional phase, Estrogen, Depression, Anxiety, Psychosocial impact, Hormonal fluctuations.*





From infancy through adulthood, women represent a dynamic interaction of biological, social, and cultural elements that influence their identities and social contributions. In terms of biology, women usually inherit two X chromosomes, which control the development of their reproductive systems and physical characteristics like wider hips and a higher body fat percentage that are linked to fertility and childbirth. Although biology provides the framework, women's roles go well beyond it. These include issues that feminist movements and policy reforms continue to address, such as systemic violence, limited access to economic and educational opportunities, and gender discrimination. Women in Tamil Nadu's districts like Coimbatore actively work in important industries like textiles, information technology, healthcare, and education, but they still have to deal with job segregation, pay inequality, and the strain of juggling work and home obligations.


Women undergo major life transitions as they approach middle adulthood (ages 35 to 65), such as the perimenopause and menopause, which are normal biological milestones despite being accompanied by emotional and hormonal changes. As women reevaluate their life goals, pursue personal development, and work to leave meaningful legacies through


caregiving, mentoring, or community involvement, this phase also corresponds with Erikson's psychological stage of generativity versus stagnation. Women can flourish if society reframes menopause as a healthy, empowering transition and supports policies like equal pay, flexible work schedules, and access to healthcare and education. This will acknowledge women's critical role in advancing progress, building resilience, and advancing sustainable development.

Psychosocial factors like culture, social support, work stress, self-image, and mental health all have an impact on the hormonal, physical, and psychological changes that occur during menopause and perimenopause. Perceptions are shaped by cultural beliefs; transitions are made easier by robust support systems and wholesome work environments. Well-being can be enhanced by coping mechanisms such as education, lifestyle modifications, mindfulness, counseling, and hormone therapy. Relationships and job performance are impacted by body image issues, decreased libido, and emotional swings, which can occasionally lead to stress or burnout. Women can manage symptoms, develop resilience, and flourish during this normal, life-changing stage with the support of supportive environments, flexible work schedules, and mental health care.

THEORETICAL FRAMEWORK







Review of Literature

Mishra et al. (2012) conducted a nationally representative cohort study involving 695 women aged 47 to 54 in England, Scotland, and Wales. They found that a minority of women, constituting 10% (n=63), experienced a severe psychological symptom profile, reaching its peak either during or in the year following menopause. Regarding vasomotor symptoms, 14% of the women (n=83) exhibited an early severe profile, characterized by a peak occurrence shortly after menopause followed by a noticeable decline. In contrast, 11% (n=67) demonstrated a late severe profile, experiencing bothersome symptoms that escalated rapidly during perimenopause and persisted at heightened levels for four years or more post-menopause. These findings shed light on the diverse temporal patterns of menopausal symptoms

experienced by women within this age range. Such insights are invaluable for developing tailored interventions and support strategies to address the varying needs of women transitioning through menopause. Mishra et al.'s study underscores the importance of considering the individualized nature of menopausal experiences and highlights the significance of providing targeted assistance to enhance women's well-being during this life stage.

Several prevalence studies, as cited by Afshari et al. (2015), Chedraui et al. (2009), Gonçalves et al. (2013), Onya and Otorkpa (2018), and Yen et al. (2009), consistently demonstrate a high prevalence of depression among women undergoing the menopausal transition. This collective body of research emphasizes the critical need to address mental health issues during this period of hormonal change.

The findings from these studies consistently reveal a significant vulnerability to depression among women in the perimenopausal and postmenopausal stages globally. This highlights a substantial public health concern and underscores the importance of targeted interventions and support systems to address the mental health challenges associated with menopause.

Gupta and Sagar (2018) and Wig and Murthy (2015) despite the global recognition of this issue, point out a significant gap in India's National Mental Health Programme and the National Mental Health Policy of 2014. Specifically, the psychiatric health of menopausal women has not been specifically addressed within these national frameworks. This discrepancy highlights the need for a more nuanced approach to mental health policy that acknowledges the unique challenges faced by menopausal women. By integrating menopausal mental health considerations into national policies and programs, India can better address the mental health needs of this vulnerable population and work towards ensuring holistic healthcare provision for women during the menopausal transition.

Krok (2015) suggests that individuals derive their sense of meaning in life from their interpretation of their livelihood. Some may find meaning through aspects such as earning a livelihood, engaging in work, or adopting a spiritual lifestyle. Alternatively, others may discover purpose in serving society, such as by managing a charity or participating in development work. Krok's perspective underscores the diverse

sources from which individuals draw meaning in their lives. While some prioritize financial success and professional accomplishments, others prioritize spiritual growth and personal development. Additionally, the desire to make a positive impact on society and contribute to the welfare of others can be a significant source of purpose for many individuals. This understanding emphasizes the importance of recognizing and respecting the varied pathways through which individuals seek fulfillment.

By acknowledging the multiplicity of sources from which people derive meaning, society can encourage a more inclusive and supportive environment. This inclusivity allows for greater exploration of different avenues for finding purpose and meaning, facilitating personal fulfillment and satisfaction for individuals.

Singh and Sivakami's (2020) subsequent investigation into Indian women navigating menopause reveals a notable lack of medical assistance during this transitional period. The study underscores a prevalent barrier where women feel inhibited from discussing crucial sexual health topics, including menstruation and menopause, with their partners. This communication gap deprives them of the anticipated social support crucial during menopause, significantly impacting their overall quality of life and life satisfaction. Consequently, these circumstances prompt introspection regarding the existential significance of their lives. In the research, Singh and Sivakami (2020) shed light on a critical aspect of women's health, highlighting the systemic challenges hindering adequate

support for women during the menopausal transition. The findings emphasize the need for improved healthcare interventions and support systems that address the unique needs of women navigating menopause in Indian society. By facilitating open communication and providing comprehensive medical assistance, healthcare providers can better empower women to manage their menopausal health effectively and enhance their overall well-being. Singh and Sivakami's study underscores the broader societal implications of inadequate support for women during menopause.

Research by Chung and van der Lippe (2020) and Sinha (2017) suggests that employed women generally experience greater levels of life satisfaction and perceive their lives as more meaningful compared to non-working counterparts. This disparity is attributed to factors such as the supportive environment and resources provided by the workplace. Chung and van der Lippe's study emphasizes the influence of workplace dynamics, indicating that positive workplace environments contribute to the enhanced well-being and fulfillment of working women. Similarly, Sinha's research underscores the importance of workplace support, indicating that it plays a pivotal role in shaping women's perceptions of their lives.

Overall, these findings highlight the significance of the workplace in promoting life satisfaction and meaning among employed women, emphasizing the need for supportive workplace policies and practices to enhance women's overall well-being.

METHOD

Statement of the Problem

The intersection of hormonal fluctuations during perimenopause and menopause with psychosocial factors significantly impacts the working women's mental health and well-being. Coping strategies and support systems are crucial in mitigating mental health challenges during this transitional phase.

Area of the Study

The area of study for the encompassed the examination of the psychosocial implications of perimenopause and menopause specifically within the context of the workforce in Coimbatore district. This included investigating how hormonal changes during perimenopause and menopause intersect with various psychosocial factors such as workplace environment, social support networks, cultural influences, and individual coping mechanisms, and how these factors collectively influence the well-being, mental health, and professional experiences of working women in Coimbatore district.

Aim and Objectives

The aim of this research is to comprehensively explore the psycho-social impact of perimenopause and menopause among working women in Coimbatore district from a social work perspective, with the goal of informing supportive interventions and policies tailored to address their specific needs.

Objectives:

1. To explore the psycho-social experiences and challenges of working women in Coimbatore district during perimenopause and menopause.
2. To investigate how hormonal fluctuations intersect with workplace environment, social support, and cultural influences among working women in Coimbatore district.
3. To assess the impact of perimenopause and menopause on mental health, well-being, and professional experiences of working women in Coimbatore district.
4. To identify coping strategies used by working women in Coimbatore district to manage psycho-social challenges during perimenopause and menopause.

Hypothesis

Working women in Coimbatore district experiencing perimenopause and menopause exhibit varying levels of psycho-social challenges, influenced by factors such as workplace environment, social support networks, and individual coping mechanisms.

Research Design

The researcher adopted an Explanatory Research Design in the study as it could involve exploring the underlying mechanisms and factors that influence the psycho-social impact of perimenopause and menopause among working women in Coimbatore district. Here the explanatory research design aims to provide deeper insights into the underlying mechanisms and factors

influencing the research topic, contributing to a more thorough understanding of the phenomena under the investigation.

Inclusion Criteria

- Women aged between 40 and 65 years.
- Currently employed or actively seeking employment in Coimbatore District.
- Experiencing menopausal symptoms or in the perimenopausal stage.
- Willingness to participate in the study and provide information about their experiences.
- Women currently undergoing treatment for menopausal symptoms that may affect their psychosocial well-being (e.g., hormone replacement therapy).

Exclusion Criteria

- Women who have undergone surgical menopause.
- Those with a history of psychiatric disorders that could confound the results.
- Individuals who are not currently employed or seeking employment in Coimbatore district.
- Women with significant health issues that could influence their psychosocial experiences unrelated to menopause.

Sampling

The researcher collected the data from the respondents using the Non-probability sampling techniques which include the Convenience Sampling and Snowball Sampling based on the research objectives, resources, and feasibility.



Convenience Sampling: This method involves selecting participants based on their availability and accessibility. The Researcher approached the women who were working in specific organizations and Institutions within Coimbatore district and were willing to participate in the study.

Snowball Sampling: In this Sampling technique, initial participants were selected based on predetermined criteria, such as being menopausal or perimenopausal working women in Coimbatore district. These participants then referred other eligible individuals they know, creating a chain referral process.

Tools of Data Collection

The researcher adopted different tools of data collection in the research study which includes questionnaire, observations, and audio recordings.

Questionnaire: A structured questionnaire was used to gather quantitative data on specific aspects of the research from the participants. The questionnaire included both the closed-ended and open-ended questions and with predefined response options, allowing for standardized data collection and statistical analysis.

Observations: Researchers directly observed the participants' behaviors, interactions, and contexts in natural settings relevant to the research study. Observations provided first hand insights into participants' behaviors, routines, and environments, capturing rich qualitative data.

Audio Recordings: Audio recordings enabled the researcher to accurately document participants' responses and interactions for later transcription and analysis, ensuring the reliability and validity of qualitative data.

Methods of Data Collection

In this research study, the case study method is employed to collect qualitative data in a comprehensive and in-depth manner and collected 6 case studies in the research. The case study method involves the in-depth examination of a specific case or cases within its real-life context, allowing researcher to explore complex phenomena and understand the intricacies of participants' experiences. Researcher used semi-structured questionnaire with participants to explore their experiences, perspectives, and challenges related to perimenopause and menopause. Observation was conducted to observe participants' behaviors, interactions, and environments in natural settings.

Researcher provided thick descriptions of the cases under study, including detailed accounts of participants' experiences, contexts, and interactions. Thick description allowed for a rich and nuanced understanding of the cases, capturing the complexity and depth of participants' experiences.

Limitations of the Study

1. Limited generalizability beyond specific cases studied.
2. Potential biases in data collection and analysis.
3. Subjective interpretations among researchers.

4. Scope constraints may overlook important factors.
5. Resource and time limitations.
6. Ethical considerations regarding participant consent and privacy.

ANALYSIS AND INTERPRETATION

CASE STUDY 1

Key Issues/Goals/Problems:

Anita, a 49 years old working woman in Coimbatore, faces a myriad of challenges associated with perimenopause. She experienced a range of symptoms that significantly impacted her work performance and confidence. Anita encounters a lack of understanding and support from both her employer and colleagues due to the prevailing stigma surrounding menopause. This lack of support exacerbated her struggles, making it difficult for her to effectively communicate the needs and access appropriate workplace accommodations to alleviate the symptoms.

Decision Criteria:

The decision-making process in addressing the respondent's situation involved several critical criteria. It's essential to evaluate the effectiveness of interventions aimed at addressing Anita's symptoms and improving overall workplace support. The feasibility of implementing accommodations within the workplace environment needs consideration. Finally, assessing the accessibility of resources for psychoeducation and advocacy is crucial in determining the most appropriate course of action.

Assumptions:

Several assumptions underpin the proposed interventions for Anita. It's assumed that the workplace accommodations for menopausal symptoms are not only feasible but also legally mandated. There is an assumption that the respondent was willing to engage in discussions with her employer and accept support from a social worker. Furthermore, it's assumed that Anita's colleagues and employer can be receptive to education and awareness efforts regarding menopause.

Content Analysis Discussion:

The content analysis delved into the respondent's symptoms and their direct impact on her work performance and confidence. It also explored the broader societal stigma surrounding menopause in the workplace, shedding light on the challenges faced by women like Anita. The analysis assessed existing policies and practices related to workplace accommodations and employee support, identifying potential areas for improvement.

Preferred Alternative with Rationale:

The preferred alternative involved facilitating open communication channels between Anita, her employer, and colleagues. This includes creating a platform for the respondent to openly discuss her menopausal symptoms and their impact on the work. Providing psychoeducation to the respondent and her workplace peers about perimenopause, its symptoms, and potential workplace accommodations is crucial. Advocating for workplace accommodations such as flexible hours and access to a quiet space for breaks aims to alleviate Anita's

symptoms and foster a more supportive work environment.

Justification/Predicted Outcome or Recommendations:

The predicted outcome of these interventions included an improved understanding and support from the respondent's employer and colleagues. This, in turn, was expected to lead to a reduction in Anita's symptoms and an enhancement of her work performance and confidence. Ultimately, the goal is to promote enhanced workplace inclusivity and support for menopausal women like the respondent.

Conclusion:

The case study underscored the significance of addressing the psycho-social impact of perimenopause in the workplace. Moving forward, future research could delve deeper into exploring the long-term effectiveness of workplace accommodations and interventions in supporting menopausal women across various occupational settings. This research could contribute valuable insights to inform policy changes and organizational practices aimed at better supporting women during this life transition.

CASE STUDY 2

Key Issues/Goals/Problems:

Priya, a 52-year-old school teacher in Coimbatore, confronts numerous challenges during her transition through menopause. Balancing her demanding workload and family responsibilities exacerbates her emotional instability and physical discomfort, leading to conflicts

within her family and a notable decline in job satisfaction.

Decision Criteria:

In Priya's case, effective interventions are crucial to addressing her symptoms and enhancing familial support. Key decision criteria include evaluating the efficacy of family therapy sessions in fostering better communication and empathy within the family unit, as well as assessing the feasibility of connecting Priya with support groups tailored for menopausal women.

Assumptions:

This case study operates under the assumption that Priya is open to engaging in family therapy sessions and is receptive to support from both her family members and a social worker. Furthermore, it assumes that over time, Priya's family members will develop an understanding of and empathy for her menopausal symptoms.

Content Analysis Discussion:

An in-depth analysis of Priya's symptoms and their impact on her familial relationships and job satisfaction is paramount. Additionally, examining the dynamics of her interactions with her family, particularly the initial challenges faced by her husband and children in comprehending her symptoms, provides valuable insights. Furthermore, evaluating the potential benefits of family therapy sessions and support group participation in alleviating Priya's symptoms and bolstering familial support is necessary.



Preferred Alternative with Rationale:

The preferred alternative entails organizing family therapy sessions to enhance communication and empathy within Priya's family. This intervention aims to cultivate a supportive environment where Priya's symptoms are acknowledged and accommodated. Moreover, facilitating Priya's connection with support groups for menopausal women offers her the opportunity to share experiences and receive additional emotional support.

Justification/Predicted Outcome or Recommendations:

Anticipated outcomes of these interventions include an amelioration of Priya's familial relationships, a reduction in conflicts, and an overall increase in support from her family members. Additionally, it is expected that Priya's job satisfaction will improve due to decreased stress and heightened emotional well-being.

Conclusion:

This case study underscores the pivotal role of familial support in mitigating the psycho-social impact of menopause. Future research endeavors may delve into investigating the long-term effects of family therapy and support group participation on women's well-being during the menopausal transition.

CASE STUDY 3

Key Issues/Goals/Problems:

Rekha, a 49-year-old factory worker, faces discrimination and isolation from her male colleagues due to her menopausal symptoms. This leads to a hostile work environment, decreased

productivity, and feelings of worthlessness, ultimately causing her to contemplate resigning from her job.

Decision Criteria:

The decision-making process involves:

- **Effectiveness of Interventions:** Evaluating interventions to address Rekha's workplace discrimination and improve her well-being.
- **Feasibility of Implementation:** Assessing the practicality of implementing sensitization workshops and legal advocacy.
- **Accessibility of Resources:** Ensuring that Rekha has access to necessary legal resources and support systems.

Assumptions:

- **Legal Protections:** Assumes that Rekha is entitled to legal protection against workplace harassment and discrimination.
- **Willingness to Engage:** Assumes Rekha's willingness to participate in interventions aimed at addressing her workplace issues.
- **Potential Attitude Change:** Assumes that Rekha's colleagues are receptive to sensitization workshops and can change their behavior.

Content Analysis Discussion:

- **Workplace Discrimination:** Analysis of the discrimination and isolation faced by Rekha in the workplace due to her menopausal symptoms.
- **Impact on Well-being:** Discussion on how workplace discrimination affects Rekha's mental health, productivity, and job satisfaction.
- **Intervention Strategies:** Examination of the social worker's advocacy efforts,



including sensitization workshops and legal support, to address workplace harassment.

Preferred Alternative with Rationale:

- Conduct Sensitization Workshops: Preferred alternative involves organizing sensitization workshops for Rekha's colleagues to educate them about menopause and foster empathy and understanding towards her situation.
- Provide Legal Support: Additionally, the social worker connects Rekha with legal resources to address workplace harassment, ensuring she has access to necessary legal protections.

Justification/Predicted Outcome or Recommendations:

- Improved Workplace Environment: Anticipated outcome includes a reduction in workplace discrimination and a more supportive environment for Rekha, leading to increased job satisfaction and productivity.
- Legal Resolution: Connection with legal resources is predicted to provide Rekha with the means to address workplace harassment and seek justice for her mistreatment.

Conclusion:

The case underscores the importance of addressing workplace discrimination against menopausal women like Rekha. By implementing sensitization workshops and providing legal support, it is possible to create a more inclusive and supportive work environment, ultimately promoting the well-being and dignity of menopausal workers.

CASE STUDY 4

Key Issues/Goals/Problems:

Kavitha, a seasoned nurse, faces several challenges as she navigates perimenopause in her demanding workplace. These challenges include unexpected emotional turmoil, a decline in empathy towards patients, conflicts with colleagues, compromised patient care, and her reluctance to seek help due to fear of judgment and isolation.

Decision Criteria:

- Effectiveness of interventions in addressing Kavitha's emotional distress and restoring her empathetic connection with patients.
- Feasibility of implementing support mechanisms within the healthcare setting to assist menopausal healthcare professionals.
- Accessibility of resources for mental health support and stress management tailored to Kavitha's needs.

Assumptions:

Assumptions include recognizing menopause as a contributing factor to Kavitha's emotional changes, assuming the availability of supportive colleagues and resources within the healthcare setting, and acknowledging the importance of addressing the psycho-social impact of menopause on the workforce.

Content Analysis Discussion:

Analyzing Kavitha's experience involves examining the emotional changes she undergoes during perimenopause, the impact on her professional performance and patient care, conflicts with colleagues, and her reluctance to seek help despite evident distress. It also involves exploring



the responses of the healthcare institution and colleagues to Kavitha's challenges.

Preferred Alternative with Rationale:

Implementing a comprehensive support program tailored to menopausal healthcare professionals like Kavitha. This includes counseling services, peer support groups, and education and awareness initiatives. The rationale is to address Kavitha's emotional needs, promote a supportive workplace culture, and improve patient care quality by restoring empathy.

Justification/Predicted Outcome or Recommendations:

The anticipated outcome of implementing the support program is improved well-being and job satisfaction

for Kavitha and other menopausal healthcare professionals, enhanced patient care quality, and greater retention of experienced staff. Recommendations include proactive measures to support menopausal healthcare professionals and foster a supportive workplace environment.

Conclusion:

Kavitha's experience underscores the need for healthcare institutions to recognize and address the psycho-social impact of menopause on their workforce. By implementing tailored support programs and fostering a supportive workplace culture, healthcare institutions can promote the well-being of their staff and improve patient care quality.

THEMATIC ANALYSIS

QUESTIONS	INITIAL CODE	THEMES GENERATED
Q. How do the common symptoms experienced during perimenopause affect your work performance, confidence, and their ability to effectively manage job duties in similar professional roles, considering the challenges they encounter?	1. Work Performance and Cognitive Impairment. 2. Emotional Regulation and Interpersonal Challenges. 3. Confidence and Job Insecurity. 4. Overall Impact of Perimenopause on Work Functioning.	Navigating Workplace challenges during perimenopause.
A. Perimenopausal symptoms significantly impact my work performance and confidence as a customer service representative. Memory lapses and		

difficulty concentrating
lead to errors and delays.
Hot flashes and irritability

hinder professional
interactions, resulting in
complaints and negative
reviews. This erodes my
confidence and raises
concerns about job
security.

Overall,
perimenopause presents
numerous
challenges,
affecting my ability to
manage job duties
effectively and impacting
my performance and
confidence
in my role.

Q. How do your
menopausal symptoms
impact the familial
relationships, job
satisfaction, and which
interventions would be
most
effective in fostering
better
communication, empathy
within your family, and
connecting you with
support groups for
menopausal women,
ultimately leading to
anticipated positive
outcomes?

A. My menopausal

1. Impact on Familial
Bonds and Job
Satisfaction.
2. Effectiveness of
Family Therapy in
Enhancing
Communication
and Empathy.
3. Benefits of
Connecting with
Menopausal
Support Groups.
4. Improvement in
Emotional
Well-being and
Conflict
Resolution

Strengthening Support
Networks for Menopausal
Well-being.



symptoms strain familial bonds and diminish job satisfaction. Effective interventions, like family therapy, foster empathy and communication within my family. Connecting with menopausal support groups provides emotional backing. These interventions helped me to alleviate conflicts, enhance my familial understanding, and to improve emotional well-being, led to a more positive outcome amidst the challenges of menopause.

Q. Could you effectively address the workplace discrimination due to menopausal symptoms, to ensure access to legal protections and to create a supportive environment for yourself and your colleagues?

A. I can effectively address workplace discrimination by holding sensitization workshops for my colleagues to educate them about menopause. Connecting with legal resources will ensure

1. Seeking Support from Peers or Mentors.
2. Practicing Self-care through Mindfulness and Exercise.
3. Participation in Stress
4. Management Workshops.
5. Adjusting Workload to Meet Changing Needs.

Coping Strategies for Perimenopausal Healthcare Professionals.



access to protections against harassment. These actions will create a supportive environment, enhancing well-being and productivity in the workplace.

Q. Why is it important for you to effectively manage her menopausal symptoms in the education sector while receiving the necessary support from school management to maintain your teaching effectiveness and to alleviate stress and anxiety?

A. Effectively managing my menopausal symptoms is crucial in the education sector to uphold teaching effectiveness and reduce stress. Support from school management is essential to validate my experiences, provide accommodations, and alleviate anxiety, fostering a conducive work environment for me and my students.

1. Managing Menopausal Symptoms Effectively.
2. Support from School Management.
3. Alleviating Stress and Anxiety.

Effective Symptom Management and Support in the Education Sector.

Q. Why is it challenging for you, as a customer

1. Menopausal Symptoms'

Supporting Menopausal Customer Service



service representative, to manage your menopausal symptoms during long working hours, and how can workplace accommodations and support alleviate these challenges to ensure job security and well-being?

A. Managing menopausal symptoms during long working hours is challenging for me as a customer service representative due to frequent hot flashes and irritability, impacting interactions with customers and job performance. Workplace accommodations like flexible schedules and support for symptom management can alleviate these challenges, ensuring job security and well-being.

- | | | |
|----|--|---|
| | Interference with Work Performance. | Representatives for Enhanced Performance. |
| 2. | Customer Interaction Challenges due to Hot Flashes and Irritability. | |
| 3. | Workplace Support Necessity for Menopausal Employees. | |
| 4. | Strategies for Maintaining Job Security and Mental Well-being. | |

FINDINGS AND SUGGESTIONS

1. Comprehensive case studies and observations in Coimbatore district reveal that working women undergoing perimenopause and menopause encounter various psycho-social challenges, including mood swings, diminished self-esteem, and increased interpersonal conflicts at work.
2. The study highlights the tangible impact of perimenopause and menopause symptoms on professional lives, such as fatigue, hot flashes, and cognitive changes leading to decreased productivity, compromised concentration, and impaired decision-making abilities.
3. Despite significant challenges, working women demonstrate resilience by employing coping mechanisms like seeking social support from colleagues,

- engaging in stress-relieving activities, and modifying work schedules.
4. Perceptions of social support among menopausal and perimenopausal working women vary, with some feeling adequately supported while others express dissatisfaction, indicating the need for targeted interventions to bolster support systems.
 5. Evaluation of existing workplace policies reveals strengths like flexible work arrangements and employee assistance programs, but gaps exist in addressing the specific needs of menopausal and perimenopausal women, highlighting the necessity for tailored policies.
 6. The study proposes actionable recommendations for social work interventions and policy development, emphasizing the creation of supportive work environments through awareness campaigns, training programs for managers and colleagues, and implementing menopause-friendly workplace policies.
 7. Addressing the psycho-social dimensions of menopause within the professional sphere is crucial, considering the challenges faced by working women during this life transition.
 8. Effective coping strategies play a vital role in navigating the psycho-social complexities associated with perimenopause and menopause, contributing to enhanced well-being in the workplace.
 9. Recognizing the importance of social support and fostering supportive work environments can empower menopausal and perimenopausal working women to thrive amidst challenges.

10. By implementing supportive workplace policies and interventions tailored to the needs of menopausal and perimenopausal women, organizations can promote their psycho-social well-being and overall professional success in the Coimbatore district.

Suggestions:

1. Develop targeted awareness campaigns to educate employers, colleagues, and working women themselves about the psycho-social impacts of perimenopause and menopause, aiming to reduce stigma and increase understanding.
2. Introduce training programs for managers and colleagues to equip them with the knowledge and skills necessary to support menopausal and perimenopausal working women effectively, fostering a more inclusive and empathetic workplace culture.
3. Establish support groups or networks specifically tailored to menopausal and perimenopausal working women within Coimbatore district, providing a platform for sharing experiences, seeking advice, and offering mutual support.
4. Advocate for the implementation of menopause-friendly workplace policies, such as flexible work arrangements, temperature control measures, and access to appropriate healthcare resources, to better accommodate the needs of this demographic.
5. Collaborate with healthcare professionals and organizations to provide accessible and comprehensive health services addressing the physical and psychological aspects of



perimenopause and menopause, ensuring working women have the support they need to manage their symptoms effectively.

6. Encourage the development of employee assistance programs that offer confidential counseling and support services tailored to the psycho-social challenges faced by menopausal and perimenopausal working women, promoting mental well-being in the workplace.
7. Conduct regular assessments and evaluations of workplace policies and support mechanisms to identify areas for improvement and ensure they remain responsive to the evolving needs of menopausal and perimenopausal employees.

CONCLUSION

The study on the psycho-social effects of perimenopause and menopause among employed women in Coimbatore district highlights the urgent need for

targeted interventions and supportive workplace policies. Findings indicate a variety of struggles experienced by women going through perimenopause and menopause, such as mood swings, decreased self-confidence, and lower productivity.

Nevertheless, the research also points out the resilience of these women and the effectiveness of coping mechanisms like seeking social support and engaging in stress-relief activities. To effectively address these challenges, there's a demand for thorough awareness campaigns, training initiatives for managers and colleagues, and the establishment of workplace policies accommodating menopausal and perimenopausal women.

By nurturing a more inclusive and supportive environment, organizations can empower these women to navigate this life transition with increased ease and well-being.



REFERENCES

- Cortés, Y. I., Marginean, V., & Berry, D. (2020). Physiologic and psychosocial changes of the menopause transition in US Latinas: a narrative review. *Climacteric*, 24(3), 214–228. <https://doi.org/10.1080/13697137.2020.1834529>
- Derry, P. S. (2004). Coping with Distress During Perimenopause. *Women & Therapy*, 27(3–4), 165–177. https://doi.org/10.1300/j015v27n03_12
- Hengel, K. M. O., Soeter, M., Maur, M., Van Oostrom, S. H., Loef, B., & Hooftman, W. (2023). Perimenopause: Symptoms, work ability and health among 4010 Dutch workers. *Maturitas*, 176, 107793. <https://doi.org/10.1016/j.maturitas.2023.107793>
- Hybholt, M. G. (2022). Psychological and social health outcomes of physical activity around menopause: A scoping review of research. *Maturitas*, 164, 88–97. <https://doi.org/10.1016/j.maturitas.2022.07.014>
- Labra, Ó., Castro, C., Wright, R., & Chamblás, I. (2020b). Thematic Analysis in Social Work: a case study. In *IntechOpen eBooks*. <https://doi.org/10.5772/intechopen.89464>
- Mendle, J., Ryan, R. M., & McKone, K. M. P. (2018). Age at Menarche, depression, and antisocial behavior in adulthood. *Pediatrics*, 141(1). <https://doi.org/10.1542/peds.2017-1703>
- Naeem, M., Ozuem, W., Howell, K. E., & Ranfagni, S. (2023). A Step-by-Step process of thematic analysis to develop a conceptual model in qualitative research. *International Journal of Qualitative Methods*, 22. <https://doi.org/10.1177/16094069231205789>
- Natarajan, J. (2013). Review Literature on Distress during the Menopausal Transition and Their Impact on the Quality Of Life of Women: What is The Evidence? *IOSR Journal of Nursing and Health Science*, 2(4), 01–10. <https://doi.org/10.9790/1959-0240110>
- National Institutes of Health State-of-the-Science Conference Statement: Management of Menopause-Related Symptoms. (2005). *Annals of Internal Medicine*, 142(12_Part_1), 1003. https://doi.org/10.7326/0003-4819-142-12_part_1-200506210-00117
- Noble, R. E. (2005). Depression in women. *Metabolism*, 54(5), 49–52. <https://doi.org/10.1016/j.metabol.2005.01.014>
- Perimenopause. (2021, August 8). Johns Hopkins Medicine. <https://www.hopkinsmedicine.org/health/conditions-and-diseases/perimenopause#:~:text=Perimenopause%20is%20the%20transitional%20time,last%202%20to%2010%20years.>
- Professional, C. C. M. (n.d.). Menopause. Cleveland Clinic. <https://my.clevelandclinic.org/health/diseases/21841-menopause>
- Sathiyaseelan, A., Patangia, B., Chaudhury, S., & Ashwini, M. (2022). How perimenopause shaped perspectives on life during the pandemic: A study on india's working and non-working women. *i-Manager's Journal on Nursing*, 12(3), 16. <https://doi.org/10.26634/jnur.12.3.19144>
- Teede, H., Misso, M., Costello, M., Dokras, A., Laven, J. S., Moran, L., Piltonen, T., Norman, R. J., Andersen, M., Azziz, R., Balen, A., Baye, E., Boyle, J., Brennan, L., Broekmans, F. J., Dabadghao, P., Devoto, L., Dewailly, D., Downes, L., . . . Yildiz, B. O. (2018). Recommendations from the international evidence-based guideline for the



assessment and management of polycystic ovary syndrome. *Clinical Endocrinology*, 89(3), 251–268. <https://doi.org/10.1111/cen.13795>

Verdonk, P., Bendien, E., & Appelman, Y. (2022). Menopause and work: A narrative literature review about menopause, work and health. *Work-a Journal of Prevention Assessment & Rehabilitation*, 72(2), 483–496. <https://doi.org/10.3233/wor-205214>